

PART ONE

1. THE STATE OF AGRICULTURAL BIODIVERSITY IN THE FARM ANIMALS SECTOR

1.1. Introduction

1.1.1. Physico-geographic and climatic features

Ethiopia is located in the horn of Africa, bordering Eritrea in the north, Djibouti and Somalia in the east, Kenya in the south and Sudan in the west. The country stretches from 3⁰N of the equator to latitude 15⁰N and from 33⁰E to 48⁰E longitude. With a land area of 110 million hectares, Ethiopia is the ninth largest country in Africa (MoA, 2004).

Ethiopia is a country of great geographic diversity. Erosion, volcanic eruptions, tectonic movements and subsidence have occurred for centuries in the country and still continue to occur accentuating the unevenness of the surface. As a result, Ethiopia is subjected to wide latitudinal and physico-geographic variations. The latitudinal variation of the country ranges from 110 meters below sea level in the Danakil depression to the highest peak of 4,620 meters above sea level (masl) on Mount Ras Dashen. The physico-geographic features are composed of high and rugged mountains, flat-topped plateaux, deep gorges, incised river valleys and rolling plains. The western and southeastern highlands are separated by the Great Rift Valley that runs from northeast to southwest of the country. Extensive semi-arid lowlands in the east, south and west are extensions of these highlands.


The Ethiopian highlands cover the central lava highlands, the southwestern plateau and the southeastern highlands. The southeastern highlands have high mountains on their western rims having slopes that run continuously towards the southeastern lowlands.

Macro- and micro-climatic conditions of the country are highly variable. The rainfall distribution is seasonal. The major rainy season lasts from April to September followed by short rainy season that occurs between September and April. The mean

annual rainfall ranges from 500 mm to 2800 mm. The southwestern regions receive the heaviest annual rainfall, which, in some areas, goes up to 2800 mm. Rainfall is moderate in the central regions and declines towards northeast and eastern Ethiopia. Annual rainfall in the southeastern and northern regions is about 700 mm and 500 mm, respectively. Similarly, temperature variations are wide. During certain seasons, temperatures go above 30⁰C or below 10⁰C in one place or another. Regimes of relative humidity, influenced by the rainfall patterns and temperature levels, are also highly variable. Because of the combined effects of the above factors, the country is endowed with diverse ecosystems that are inhabited by amazingly great diversity of animal, plant and microbial genetic resources (FAO, 2001).

Ethiopia is comprised of nine federal states and two city councils (Figure 1). Eighty-three distinct languages having 200 dialects are spoken. Amharic is the working language of the Federal government. As per the provision of the Constitution, official languages used in the regional states are those that are chosen by the respective states. English is widely used in business and academic circles (MoA, 2004).

Figure 1. Regional States of the Federal Democratic Republic of Ethiopia



1.1.2. Population

The Estimated human population of Ethiopia in 1980 and 1990 were 37.7 million and 50.8 million, respectively with an average annual growth rate of about 3%. Today, Ethiopia is the third most populous country in Africa. The estimated current population is 69.1 million with annual rate of growth at 2.7% (Annex Table 1.7). The projected human population is estimated at 84, 106 and 129 million in 2010, 2020 and 2030, respectively. Females constitute 49.8% of the current population. Forty-seven percent of the females, i.e. 24% of the population, are of childbearing age. The population between ages 15 and 59 years constitutes 52.2% of the current total, while the population under 15 years constitutes about 43%. The level of fertility, at about 5.9 during the current year, is among the highest in sub-Saharan Africa. The level of total fertility in rural areas is higher than in the urban areas. According to the population census of 1994, for example, total fertility of the rural areas (6.4) was almost twice as high as in the urban areas (3.3). Crude birth and death rates (number of births and deaths per 1000 population per annum) between 1994 and 2000 were 44.4 and 44.2, and 16.4 and 15.0, respectively. In 1994, life expectancy of females and males was about 52.7 and 50.2 years, respectively (Genet Mengistu, 2004).

The majority of Ethiopians dwell in rural areas. The proportion of urban dwellers until about 1940s was less than 3%. Over the years, this proportion increased reaching 9.7, 11.4, 13.8 and 15.0% in 1970, 1984, 1994 and 2004, respectively (Annex Table 1.7). According to projections made by Genet Mengistu (2004) the proportion of the urban dwellers in 2010 and 2020 will be 17.2% and 19.9%, respectively.

Distribution of the population is highly skewed towards the highlands. This is attributed mainly to different environmental factors. About 80% of the population is settled in 45% (above 1500 masl) of the country's land area. As a result, the highland areas are more densely populated than the lowland areas, which are characterized, by insufficient rainfall, high temperature regimes and vector infestation such as by tsetse flies. The highland areas, therefore, have for long been subjected to severe overgrazing and degradation of its natural resources.

1.1.3. Agriculture

Agriculture is the main stay of the Ethiopian economy employing about 85% of the total population. It contributes about 45% to the GDP, 90% to the total export earnings and 70% of the raw materials to the agro-industrial sector. Smallholders still dominate at present as they generate more than 95% of the overall national agricultural output. Despite this, due to limited access of the smallholders to agricultural inputs and extension services, farming is still practised in traditional ways (MoA, 2004). According to CSA (2004), for example, the proportion of the total cultivated land where fertilizer, pesticides and improved seeds are applied/used constitutes only 36.8, 8.2 and 2.8%, respectively. Similarly, the proportion of the crop-farming households (totalling about 11 million) who participate in the extension package is only 5.6%. The level of education of the farming community is very low. Illiterate farming households comprise about 64.6% of the total.

About 66% of the country's land area is suitable for agriculture but only 22% of this area is currently under cultivation. Similarly, only 4.4% (0.16 million ha) of the country's potentially irrigable land is under irrigation (MoA, 2004), and about 90% of total cultivated area is occupied by crops. More than half the country's land mass is suitable for grazing and browsing (Annex Table 1.2). Land tenure for livestock production is under private and communal grazing (Annex Table 1.4).

Farm animals are an integral part of the Ethiopian agricultural fabric. They are sources of food, traction, manure, raw materials, investment, cash, security, foreign exchange earnings, social and cultural identity. The success of Ethiopian agriculture, in addition to other factors, is based on availability and/or contribution of farm animals and no agricultural practices other than mowing are conducted without the involvement of farm animals. Land cultivation is mainly done by oxen, horses, donkeys or their combinations. Farm animals serve to level the ploughed field, shortly before and after sowing. In some areas, farm animals are involved in weeding, especially maize and sorghum fields. Transportation of the harvested crops to and from threshing sites, threshing itself, transportation to and from the market is conducted by the farm animals. Similarly, transportation of water, firewood, mobile houses, construction

materials and other goods is conducted by farm animals and they are the main means of human transport.

Livestock have several other values that are decisively important for the livelihood of the farming communities. Their primary products such as meat, milk, egg and honey are major parts of the staple diets and prominent sources of animal protein. Live animals, skins and hides are sources of cash income. Moreover, skins, hides and horns are used as construction materials for a range of household items, agricultural tools and ornaments. Their manure is used to fertilize crop fields. Farm animals serve as insurance and a source of cash for rural farming communities. They are profitable areas of investment. Their role in cultural and social ceremonies is significant. Major primary products of the livestock are presented in Annex Table 1.8.

Animal dung is used as a construction material. It is also the major source of fuel, particularly in areas where firewood is scarce. According to NCS Secretariat and MNRDEP (1994), the contribution of manure to the total energy consumed in 1984 was about 7.5%. In 1990/91, about 3.9 million tons of dung was consumed. According to the above source, the increase in the annual demand for traditional energy and proportion of dung that was used as fuel in the early '90s was estimated at 1.5 million tons and 70%, respectively.

Livestock play a decisive role in achieving food self-sufficiency in the country. The livestock sub-sector is one of the input suppliers to the manufacturing sector of the country. According to Sileshi Ashine (2003), the contribution of livestock to the total agricultural GDP and national foreign currency earnings is about 30% and 16%, respectively (Annex Table 1.1). Major export items from the sub sector comprise live animals, hides, skins, frozen meat, dairy products, honey and beeswax (Annex Table 1.10).

Given the country's diverse topographic and climatic conditions, a huge livestock population, different breeds of animals evolved over time and adapted to the ecological conditions of their habitat; and to some extent been influenced by production systems,

Ethiopia could be considered as a centre of diversity of animal genetic resources (Beyene Kebede and Beruk Yemane, 1992).

Despite, *inter alia*, the uncompromising contributions the livestock sub sector plays in the livelihoods of the majority of Ethiopians and in the overall national economic development, the attention given to develop the sub sector has not been so significant. Needless to say, this is despite their huge diversity and intrinsic capability to adapt to adverse and ever-worsening environmental conditions and the future implications of this potential.

Thus, the fate of production and productivity of the sub sector is still left to depend on the scarce and ever-declining marginal and commonly used lands. Provision of health facilities and services are far below the required minimum. They have been left to undergo untraceable levels of inbreeding and crossbreeding. And yet, depicted as the "low performers", they are expected to gradually give way to other "best performing" genotypes. Consequently, if indiscriminate distribution of the "best performing exotic genotypes" to different parts of the country is to continue at the current pace, the gene pool of the locals, particularly of the chicken, would be lost in the near future before they are even fully described and understood.

Major farm animal species of the country are cattle, sheep, goats, horses, mules, chicken and camels (Annex Table 1.6). Honeybees are often considered as farm animals. Diversity and composition of species of livestock are variable from place to place. This is mainly due to differences in agro climatic conditions. At lower altitudes, the proportion of goats and camels rises. The majority of livestock are in the rural areas, but limited numbers of some species are also kept in the peri-urban and urban areas.

1.2. Distribution and Roles of Livestock in Major Production Systems

1.2.1. Mixed crop-livestock system

Mixed farming system is predominantly found in highland agro-ecological zones where the climatic factors are conducive for farming of crops and raising livestock. In this system, livestock and crops are maintained as complementary enterprises. The average land size per household is often less than two hectares. Consequently, the intensity of farming practices often moderate to high albeit the production is mainly subsistence in nature. The relative importance of livestock products and services within species has not been properly quantified. Thus, Annex Tables 4.1 and 4.2 could not be completed.

The major species of farm animals of the country, except camels, are found in this farming system. Although dominated by cereal crops production, the diet of the people is composed of cereals, vegetables, meat and milk.

Compared to other farming systems, this system is more conducive to crossbred dairy cattle production. The system is characterized by land scarcity, severe resources degradation and recurrent drought. Former productive grazing lands are gradually turning into crop fields. Therefore, major feed sources are grazing marginal lands, crop aftermath and crop residues. Crop residues are the major sources of feed, particularly during periods of feed shortage. In the rural areas, dung is the major source of fuel.

1.2.2. Agro-pastoral system

In the agro-pastoral system, human pressure on natural resources is relatively lighter than that observed in higher altitudes. Household landholding is often greater than in the mixed farming system. Livestock are important components of the farming system. Crops are produced both for subsistence and market. Livestock are kept for draft, sale and generation of other primary products. All of the major species of Ethiopian farm animals are possibly found in this system, but with variable species composition. The lower the altitude the higher will be the proportion of small ruminants, especially goats. Livestock are mainly kept in communal grazing lands and the use of crop residues and aftermath grazing is not uncommon. The diet of the people is composed

of cereals, vegetables, meat and milk. The use of dung for fuel is not as intense as in the mixed crop-livestock production system.

1.2.3. Pastoral system

Pastoralism is practised in vast arid agro-ecological zones of Afar, Somali and Borena rangelands. Land ownership at household level is not a common practice. Areas like the Somali rangelands are mostly held by different clans. Despite their vast size, pastoral areas are sparsely populated compared to the other farming systems. As a result, they are subject to intrusions from the highlanders and large-scale commercial plantation schemes. Crop production is not a feature of the system and subsistence is almost entirely based on livestock and livestock products. The main source of food is milk. Consequently, pastoralists tend to keep large herds to ensure sufficient milk supply and income. Although most of the farm animal species, excepting horses and mules, are reared in this system, it is dominated by goats, cattle, sheep and camels.

Species of indigenous farm animals found in this farming system, except for insignificant cases in chicken, is not diluted by exotic blood. Distribution of exotic chicken is still being promoted and carried out. Water and feed shortage are the most limiting factors of production and productivity of the system.

Water development activities, if available, are limited mainly to accessible areas. Entire sources of feeds are rangelands. Herdsmen have virtually no control on their productivity. Therefore, pastoralists are forced to move constantly to different areas, sometimes even into the neighbouring countries in search of feed and water for their animals. Cattle are often moved out of their area during such times. Camels are used to transport mobile houses of the herdsmen. Such movements are considered the life-saving strategies adopted for decades. However, they have also some undesired effects such as triggering of conflicts between the occupants of the areas and the newcomers. They also create the opportunity to the mixing up, around potential watering and grazing/browsing areas, of formerly separated herds of different stocks and species that came from various areas. This results in interbreeding. In addition to several other factors, gradually increasing human and livestock populations are further deteriorating

the productivity of rangelands and severe droughts frequently occurring in the area are worsening the situation. Moreover, some of the productive rangelands are being invaded at an alarming rate by exotic weeds such as *Prosopis juliflora*.

1.2.4. Peri-urban and urban system

Practically all species of farm animals available in the country are kept in peri-urban and urban areas. In recent years, peri-urban and urban commercial dairy and chicken (both layers and broilers) production is increasing in and around cities and major towns. The level of management adopted currently in peri-urban and urban livestock production system varies from relatively low to high intensive systems (Azage et al., 2001).

The expansion of peri-urban and urban livestock production system, particularly commercial dairying and chicken rearing, is favoured by multitudes of opportunities. Like any other emerging economic system, however, it is yet to be tested against challenges emerging simultaneously in such areas as management, disease control, reproduction, quality control and public health, waste management and disposal, and its ultimate success will depend on how best the system could overcome these challenges.

1.3. Levels of Inputs in Different Production Systems

Based on the type and quantity of inputs as well as the objective and level of intensity of production, major livestock production systems presented above could be further classified into low, medium and high input level production systems (Annex Table 2.1).

The level of intensification of the systems depends on land tenure, social structure, know-how, access to credit and inputs, availability of market and economic return. Similarly, the level of utilisation of inputs depends on climatic conditions, production systems, types of livestock and livestock products demanded, species and breed of animals used, the extent of use of crossbreeds, accessibility to consumer areas,

availability of reliable sources of inputs, prices of inputs, the level of know-how, availability of labour, pasture and other feed sources, and the level of extension services rendered.

1.3.1. Low input production system

Low input production system is a production system where one or more rate-limiting inputs impose continuous or variable severe pressure on livestock, resulting in low survival, reproductive rate or output. Output and production risks are exposed to major influences, which may go beyond human management capacity (FAO, 2000).

The majority of the country's current livestock production system lies in the low input category. It accommodates more than 95% of the livestock population. The main features of the low input production system are its full dependence on natural resources and the limited demand for inputs. Low input production system is found in three of the four livestock production systems prevailing in the country, namely: pastoral, agro-pastoral and livestock-crop mixed farming systems. However, it is more pronounced in pastoral livestock production system than in the others (Annex Tables 2.4, 2.5, 2.6, 2.7, 2.8 and 2.10).

Most of the cattle in low input production system are indigenous breeds. Considerable numbers of crossbred dairy cows are also distributed into these areas. Based on agro-ecology and tradition, the uses of cattle in the system are variable.

In the lowland areas, cattle are used mainly for milk production. To ensure a constant supply of milk, therefore, a large number of females and few selected breeding males are maintained. The remaining males are sold while young. These areas provide about 20% of the draft animals used in the highland areas. About 90% of live animals exported are produced from these areas (Sileshi Ashine, 2003). Moreover, the areas supply considerable number of cattle to those who are involved in fattening and crossbreeding practised in smallholder dairy programs. The lowland areas are, thus, important sources of livestock supply to the neighbouring highlanders.

In the mixed crop livestock farming dominated highland areas, cattle are mainly reared for draft purposes. Milk and meat are considered as secondary priorities. Fattening or conditioning of oxen and/or steers is practised in the central highlands. The practice is aimed at selling with relatively good prices of old draft oxen after feeding for a specific period of time. Young replacement oxen and steers are purchased at cheaper prices.

In the low input production system of the highlands, small ruminants are the second most important animals next to cattle. They are prominent sources of meat, skin, milk, wool/hair and cash income. This is true for the lowland areas also, except that they are not used for wool/hair production. Except for a few crossbred animals resulting from the distribution of imported rams, the majority of the small ruminants are indigenous breeds. Most of the sheep population is found in the highland areas while goats are almost equally distributed in the highland and lowland areas (FARM Africa, 1996).

Most of the equine populations of the country are found in the low input production system and are all indigenous species. They are the main means of transportation for humans and goods. In some parts of the country, especially in the highland areas, horses and occasionally donkeys are also used for ploughing, threshing and other agricultural activities.

The entire camel population of the country is maintained in the low input production system of the lowland areas. All camels are indigenous breeds and provide milk, meat, transport and other sources of income as well. Camel milk constitutes the major part or even the entire part of the diets, thus serving as the backbone for the livelihoods of some pastoralists (Coppock, 1994).

More than 99% of the poultry population is found under the low input production system. Almost all the poultry are local breeds. Very few improved and crossbred poultry are found under this system. In low input production system, poultry is one of the major sources of food and cash income.

1.3.2. Medium input production system

Medium input production system is a production system where management of the available resources has the scope to overcome the negative effects of the environment, although it is common for one or more factors to limit output, survival or reproduction in a serious fashion (FAO, 2000).

Medium-input production system consists of farmers who are using relatively different kinds of inputs. The majority of these farmers are residents of urban and peri-urban areas. About 1% of the cattle and 4% of the chicken population are found in this production system. The smallholders participating in the government livestock extension package programs are heading towards the medium-input production system. Moreover, the expansion and the success of this system are favoured by numerous opportunities.

Species with good market opportunities are involved in the medium input production system. As presented in Annex Tables 2.3 and 2.9, only cattle and poultry are treated under the system.

Data on change in the distribution of production systems over the last 20 years (which was supposed to be filled in Annex Table 2.2) are not available. Thus, further study is required to illustrate the distribution over time.

1.3.3. High input production system

High input production system is a production system where all rate-limiting inputs to animal production can be managed to ensure high levels of animal survival, reproduction and output. Output is constrained by managerial decisions (FAO, 2000).

Although the trend is towards high input production and there are few private poultry, dairy and fattening farms, the system is at the moment not significant.

1.4. The State of Farm Animal Genetic Resources Diversity

In Ethiopia, classification of farm AnGRs into breeds is far from complete. Classification studies have been conducted on most of the cattle and goat breeds that exist in the country. However, research is at its rudimentary stage for the other species, particularly chickens. Therefore, the list of breeds presented below should be viewed from this perspective.

The most common farm animals of the country can be categorized into mammalian, avian and honeybee species. Cattle, sheep, goats, camels, donkeys, horses and mules are the major farm animals that lie under the mammalian category. The number of breeds of cattle, sheep, goat, camel, donkey, horse, mule and chickens breeds identified so far are 25, 13, 15, 4, 4, 2, 2 and 5, respectively (Annex Table 3.1). Under the avian category are chicken (poultry), ostrich and turkey. The latter two avian species are not widely used in the country. Stinging honeybee species are economically the most important species in the country.

1.4.1. The state of diversity of farm AnGR

Cattle breeds

Indigenous breeds: major cattle breeds identified so far are Arsi, Begayit, Ogaden, Borena, Goffa, Arado, Nuer, Gurage, Jidu, Karayu/Afar, Harar, Horro, Smada, Fogera, Mursi, Raya-Azebo, Adwa, Jem-Jem, Sheko, Ambo, Jijiga, Bale, Hammer, Medenece and Abergelle. Medenece and Abergelle are recently reported by the Tigray Regional Bureau of Agriculture and Natural Resources Development (1999) to exist in that part of the country. In this regard, it is plausible to believe that Ethiopia may be endowed with more cattle breeds, which have not yet been described.

Out of 25 indigenous cattle breeds, the Borena, Horro, Fogera, Karayu, Arsi and Nuer are the widely used breeds (Annex Table 4.3).

Exotic breeds: The purpose of importation of exotics was to improve milk production in the country. The breeds so imported are Holstein-Friesian, Jersey and Simmental. The breeds are being used in medium input production system but the lion's share is occupied by the Holstein-Friesian and their crosses. Crossbreeds used under medium input production system are those produced from crossings between exotic sire breeds and three indigenous dam breeds, namely: Borena, Fogera and Arsi. The above crossbreeds and others resulting from crossings between Borena with Simmental, Arsi with Jersey, Horro with Holstein Friesian, Horro with Jersey, Horro with Simmental are being used by research and/or teaching institutions.

Sheep breeds

Indigenous sheep breeds: Major sheep breeds found in Ethiopia are Begayit, Farta, Horro, Abergelle, Menz, Begi-Degu, Arsi, Ille, Tukur, Bonga, Afar, Dangila and Black Head Somali (formally known as Black Head Ogaden) sheep breeds. Farta sheep breed (found in South Gondar Zone of Amhara Regional State), and Abergelle, Begi-Degu and Ille sheep breeds (found in Tigray Regional State) have been reported recently by the agricultural bureaux of respective regions.

Exotic sheep breeds: Exotic sheep breeds introduced for their wool and mutton production are Awassi, Hampshire, Blue-de-main, Merino, Romney, Corriedale and Dorper. Crossbreeding of the Menz breed with the five exotic breeds, namely: Awassi, Hampshire, Bleu-de-Main, Romney and Dorper are being used for development and research activities.

Goat breeds

Indigenous goat breeds: Major goat breeds existing in the country are Begayit, Ille, Afar, Hararghe Highland, Arsi-Bale, Short-eared Somali, Woyito-Guji, Long-eared Somali, Central Highland, Abergelle, Western Highland, Widar, Western Lowlands, Maefur and Keffa. Moreover, Felata, Arab, Gumuz, Agew and Oromo sub-types of the western lowlands have been recently reported.

Exotic goat breeds: The aim of introducing exotic goat breeds was to improve milk production of the local goat breeds. Anglo-Nubian and Toggenberg are exotic goat breeds that were introduced by Farm-Africa and higher learning institutions. Thus, crossbreeds between Anglo-Nubian and Hararghe Highland and Anglo-Nubian and Somali are being used for milk production by smallholders in central, eastern, southeastern, and southern parts of the country. Toggenberg and their crosses with Hararghe Highland are used for research purposes at the Alemaya University of Agriculture and Debub University.

Equine breeds

Donkey breeds that exist in the country are the Jimma, Abyssinian, Ogaden and Sinnar. Major breeds of horses that have so far been well recognized are the Oromo and Dongola. In Ethiopia, crossing of Asses with mares to produce mules dates back to centuries. Except for two well known, namely: Sinnar and Wollo Mule breeds, there are no other well-defined hybrids in the country.

Camel breeds

Attempt to classify Ethiopian camels has not been satisfactory so far. Wilson (1984) has classified and described major camel breeds in the country as the Afar, Borena, Anfi and Somali/Ogaden breeds.

Poultry breeds

Indigenous chicken breeds: Indigenous chicken breeds identified so far are Horro, Jarso, Tililli, Tepi and Cheffe breeds that are found in the central highland areas (Tadelle et al., 2003), and the naked-neck breed found in northern, northwestern, western and southern lowland areas of the country.

Exotic chicken breeds: Several layer, broiler and dual-purpose exotic chicken breeds introduced into the country are being used for food and agriculture. Rhode Island Red (RIR), White Leghorn, Lawman Brown, Cobb-500, Fayoumi, Bovans Brown and Arob

Acre are reared by small and large-scale commercial producers in urban and peri-urban areas. Besides, RIR and White Leghorns as well as their crosses with indigenous chicken are used by rural smallholders for egg and meat production.

Honeybees

It is estimated that Ethiopia has about 10 million honeybee colonies. Species of honeybees identified so far are *Apis mellifera adansol*, *Apis mellifera lementica*, *Apis mellifera monticola*, *Apis mellifera litorea* and *Apis mellifera abyssinica*.

1.4.2. Population trends

Decreasing

Cattle: Complete information on trends of populations of cattle at breed level is not available. Limited information available in literature and obtained from field surveys indicate, however, that there are indigenous cattle breeds with decreasing trends. These are the Sheko, Fogera, Begayit and Borena.

Sheko breed: Sheko cattle breed is the only remaining representative of the brachyceros type in eastern Africa. The total population of the breed reported by FAO (1999) was 31,000. According to the information obtained from Bench-Maji Zonal Agricultural Development Office in 2001 (the Zone in which the breed exists), however, the total population of the breed was estimated at 18,307 heads. According to the latter source, the majority of the heads so counted as Sheko breed are, unfortunately, those animals that might have been produced by interbreeding with the local zebu found in their surroundings. The major causes of decline in population are trypanosomiasis, inbreeding and interbreeding with other local breeds of cattle. It is generally believed that the breed is at risk of being lost.

Fogera breed: The Fogera is exhibiting a decreasing trend. According to information from the Amhara Regional Agricultural Development Bureau (2001), the estimated population of this breed was 800,000 in 1981 and 636,000 in 1998. The major causes

contributing to the declining trend are inbreeding, uncontrolled inter- and crossbreeding, diseases and drought.

Begayit breed: Compared to several others, Begayit cattle breed has small population size and is confined in a smaller area in the northern part of the country. Even though exact information on the population trend of the breed is lacking, it is believed that the population of Begayit cattle has suffered greatly due to high off-take rate during the Ethio-Eritrean war that occurred between 1998 and 2000.

Borena, Afar and Ogaden breeds: Complete information on the population trend is lacking for these breeds, too. Because of prevailing feed shortage resulting from bush encroachment, recurrent drought, and interbreeding and diseases; these areas are experiencing vivid change in species composition from cattle to camels and small ruminants. Therefore, cumulative effects of the above factors are believed to have negatively affected the population trend of these breeds.

In Ethiopia, artificial insemination of indigenous cattle using the semen from exotic cattle breeds has been implemented over the last three to four decades. While artificial insemination has, in most cases, been and is still being executed on indigenous breeds that have not been evaluated and/or not characterized, simultaneous efforts to conserve the gene pool of the indigenous breeds are non-existent. Due to the ease of artificial insemination, therefore, indiscriminate crossbreeding is resulting in unforeseen levels of dilution of the indigenous gene pool.

Chickens: Complete information on the current status of indigenous chicken breeds is not available. As a matter of fact, introduction of exotic breeds of chicken into different parts of the country have been conducted over the last five decades and the trend is increasing in almost all parts of the country. Such massive introduction of exotic blood into different corners of the country is conducted via distribution of fertile eggs, day-old chickens, crossbred pullets and exotic cockerels. A number of governmental and non-governmental organizations are involved in this operation. To this end, about seven poultry breeds multiplication centres are involved in the

production and distribution of exotic and crossbred chickens for pure and crossbreeding purposes aimed at improving the genetic potential of the local breeds for egg and meat production. As a result, distribution of "best performing genotypes" is being implemented at an alarming rate and if this trend continues at the current pace, the gene pool of the local breeds could be lost in the near future before they are even described.

Other farm AnGR:

Information on the trends of other farm AnGR is lacking. However, the Afar sheep and Sinnar donkey breeds are reported as decreasing.

Increasing

The only species of farm AnGR reported as increasing is the camel. This is mainly attributed to bush encroachment that particularly resulted in cattle populations decreasing.

Stable

Farm animal breeds whose population is reported as stable are the Horro cattle and sheep breeds, all indigenous goat breeds and the Black Headed Somali sheep breed.

1.5. The State of Development and Use of AnGR

1.5.1. Livestock development

This part presents the number of breeds that are actively used for food and agriculture. Where available, information on development efforts as well as information on policy and strategic issues on development and use of farm AnGR on species basis is presented.

Cattle

Although Ethiopia has no standing breeding policy, crossbreeding activities are nonetheless being conducted on four widely used cattle breeds, namely: Borena, Horro, Arsi, and Fogera (Annex Table 4.3). Borena, Arsi and Fogera are crossbred with Holstein-Friesian while Horro is crossbred with Jersey. To some extent, selection on Borena, Fogera and Horro breeds is being undertaken in different ranches (Annex Table 4.4).

Recognizing the contribution of the livestock sub sector to the national economic development and improvement of nutritional status of the people, quite a number of programs and projects have been designed and implemented. The pioneer project that started extension work on improving cattle under smallholder production system was implemented by the Chilalo Agricultural Development Unit from 1967 up to 1975 and the Arsi Rural Development Unit from 1975 up to 1984. Following these, a number of programs and projects that mainly focused on smallholders were implemented. These include:

- The Joint Program for Eradication of Rinderpest and CBPP Campaign (1969-1975),
- The Wolayita Agricultural Development Project (1974),
- The Minimum Package Program (1972-1980),
- The Second Livestock Development Project (1973-1980),
- The Third Livestock Development Project (1977-1993),
- The Fourth Livestock Development Project (1987-1991),
- Addis Ababa Dairy Development Project (1972-1981),
- Dairy Rehabilitation and Development Project (1986-1990),
- Pan Africa Rinderpest Campaign (1993-1998),
- Hides and Skin Improvement Project (1991-1997),
- Selale Peasants Dairy Development Pilot Project (1981-1994),
- Support to National Artificial Insemination Centre (1987-1980),
- Smallholder Dairy Development Pilot Project (1981-1994),

- Smallholder Dairy Development Project (1995-2000).

On-going projects that are currently contributing to AnGR are the National Livestock Development Project (1999), Integrated Livestock Development Project (1998), Pan African Control of Epizootics (2000), Farming in Tsetse Controlled Areas (2001) and Livestock Development Master Plan Study (starting from 2004).

Along these projects and programs, the regular livestock extension program, with special emphasis on dairy extension service has been implemented at all levels of the MoA structure over the last three to four decades. The extension service was delivered through various methods, such as training to experts and farmers, distribution of in-calf crossbred heifers, distribution of breeding bulls and improved forage seeds as well as delivery of artificial insemination (AI) and animal health services.

In Ethiopia, work on the development of animal genetic resources, particularly in the dairy sector, goes back to post-World War II years. Ethiopia got the first batch of dairy cattle through United Nations Relief and Rehabilitation Administration (UNRRA), under the Marshal Plan set to rehabilitate the war affected allied force countries (Solomon Haile Mariam, 1981). Since then, agricultural training institutions were established such as the Alemaya College of Agriculture and Holetta Research Station which contributed to manpower development in the field and the initiation of research in the livestock sector, particularly in dairy and extension services.

All dairy development endeavours require inputs such as improved dairy cattle, AI and improved forage seeds. Accordingly, the government established five cattle improvement and multiplication ranches from which crossbred in-calf heifers and crossbred breeding bulls are distributed to smallholder farmers. Nevertheless, there is still a serious shortage of improved dairy cattle in the country. The country used to import pure and crossbred dairy cattle for breeding and research purposes mainly from Kenya.

The efforts made so far by different projects and extension programs have helped gear smallholder farmers towards management of improved dairy cattle, improved nutrition and increased income at the household level. The dairy sector has also contributed to crop production through provision of additional power for farming, manure, and generating income for purchasing inputs needed for crop production.

Similar to the dairy sector, various governmental and non-governmental organizations have undertaken a number of livestock development and related projects. In this respect, the MoA is the principal organization followed by the Ministry of State Farms Development and the Ministry of Trade. Research activities on beef have been carried out by the Institute of Agricultural Research, Alemaya University of Agriculture, the junior agricultural colleges and other development organizations.

Cattle fattening on a small-scale is emerging as a profitable, low-risk and low-input livestock enterprise for smallholders (EARO, 2002). Studies conducted on beef fattening at Jirma State Farm and on economics of feeding old oxen for beef production at Adami Tullu indicated that beef production is a profitable enterprise. However, the livestock marketing system is not well organized in all aspects to benefit smallholders. Poor market information system, low level of organization or institutional set-up and lack of proper transportation facilities required for local and foreign markets have influenced the sector. Lack of proper understanding of the production system, inadequate nutrition, fragile environment, prevalence of diseases as well as problems associated with processing, marketing and land use, and thermal stress are other important constraints to beef production.

Major constraints and challenges to the development of the dairy sector:

- Lack of co-ordinated activities engaged in the development, research, training, extension and delivery of inputs.
- Development efforts have not been institutionalised and have rather been short-term project-oriented undertakings. They were often fragmented and were based on piecemeal approaches with no inter-linkages.

- An accountable and responsible body for the coordination and development of the dairy industry has not been identified and set in place. As a result, improved performance in husbandry and management has been hampered.

Sheep

Sheep have social and economic importance to the producers who keep indigenous breeds for meat, hair production and income generation. The sheep have coarse fleece and a coat of short or long hair. To improve their productivity, the MoA has drafted two development strategies (1) to improve and select a program for indigenous sheep with higher potential, and (2) to crossbreed small- to medium-sized local breeds with the exotics where there is no other choice for improvement.

In 1981, ARDU started a breeding program on indigenous breeds which involved (1) crossbreeding exotic with indigenous sheep to study the performance of different levels of crossing and (2) producing crossbred rams to be distributed to farmers. The indigenous breeds involved in this program were Arsi, Bonga, Horro, Black Head Somali and Afar. Rams of exotic breeds involved in the program were Bleu-de-Main, Corriedale, Romney and Hampshire (Annex Table 4.3).

MoA also established three sheep breeding and multiplication ranches in different parts of the country. The main objectives of these ranches were to improve meat and wool production of the indigenous sheep through selection and crossbreeding. The indigenous breeds that were involved under these programs were Black Head Somali, Menz and Horro.

Research activities were carried out by the former Institution of Agricultural Research (IAR), higher teaching institutions like Alemaya University of Agriculture (AUA), the Ambo and Jimma Junior Agricultural Colleges and ARDU. Most of the research activities focused on performance testing of crossbred sheep at different blood levels and to some extent breed evaluation of the indigenous Arsi, Horro, Black Head Somali and Afar sheep breeds.

Goats

Little attention has been paid to the development of goats despite their economic importance, particularly to the most vulnerable groups of the community. In 1988, FARM Africa initiated a dairy goat development program in collaboration with the MoA, AUA, Awassa College of Agriculture (ACA) and several international and local NGOs. The objectives were to identify and characterize the indigenous goats, to describe the traditional goat husbandry practices in different production systems to develop and test methods for the rapid survey of livestock.

Poultry

Chickens are widely kept in Ethiopia and make up the largest population of farm AnGRs of the country. According to Teketel Forssido (1986), the distribution of chicken varies with altitude. The local chickens (non-descriptive types) vary widely in body size, conformation, plumage and other characteristics. In depth studies to determine variations that exist in egg and meat production and the productivity among different poultry breeds have never been commissioned in Ethiopia (Million Tadesse et al., 2003). Both crossbreeds and pure exotic breeds are widely used to increase meat and egg production. A number of governmental and non-governmental organizations have been involved in distributing pure exotic breed pullets, cockerels and fertile eggs to farmers from abroad and poultry breeding and multiplication centres existing in the country. Exotic breeds used as pure and for crossbreeding with the locals are the White Leghorn, Rhode Island Red, Bovans Brown and Bubcocks (Annex Table 4.3).

Equines

Equines have been playing an important role in the country by supplying vital power for traction, transport and field operations. Animal power is the backbone of food production and rural economy of smallholders in the highlands of the country. In these areas, over 90% of the farmers use animal traction for food crop production. The density of equine population in the country is estimated to be 6.1/km². There is one equine for every five persons of the total population and for every four persons that are

engaged in agriculture. Donkeys, horses and mules together represent 13.7% of the total live weight of domestic herbivores of the country. An estimated 75% of the farms are more than a day and half walk from all weather roads. Hence, the very low proportion of villages that can be reached by all-weather roads makes delivery of goods extremely difficult. Equines are, thus, important animals in Ethiopian agriculture and in the overall national economy. (Asamenew et al. 1993).

Camels

In Ethiopia, only the one-humped camel (*Camelus dromedaries*) exists. Breeds of camels identified so far are Afar, Borena, Anfi and Somali/Ogaden breeds. There exists, however, no conceptual classification of camel breeds or strains that helps to categorize according to their potential for milk, meat and draught. Although the Faculty of Veterinary Medicine of Addis Ababa University has made several studies on diseases of camels (Bekele et al., 1998) only recently has research on camel husbandry been initiated at Alemaya University of Agriculture, now followed by Mekele University College.

1.5.2. Livestock policy and strategy

Although the country does not have a breeding policy on farm AnGR, the Rural Development Policy gives due emphasis to the agricultural sector. The policy envisages building the capacity of farmers in various ways and encourages formation of cooperatives. This has positive effect on livestock development since the majority of Ethiopian farmers are involved in livestock production. In addition, there are other sectoral policies such as the Investment Policy that has created favourable environment for investments in the livestock sub sector. This is mainly aimed at promoting labour-based strategy and broad based internal and external market opportunities.

To support the development of the livestock sub sector, the MoA has drafted a Ruminant Livestock Development Strategy in 1995. The general objective of the draft strategy is to develop and utilize the available resources and to increase its contribution to the social and economic development of the country. The other strategy, having a

sectoral nature, is the Industrial Development Strategy, which is based on Agricultural Development Led Industry. It is strongly believed that this strategy supports all agricultural sectors through creating markets for local agricultural products.

1.6. The State of Conservation of Farm AnGRs

In developing countries, animal genetic resources are being eroded through rapid transformation of agricultural systems. One of the major causes to the loss of indigenous breeds of farm AnGR is indiscriminate use of exotic genetic resources. According to FAO (1999), about 30% of the world's farm animal breeds are subjected to the risk of extinction.

Conservation of local breeds of farm AnGR is part of animal husbandry and should, ideally, be based on complete information on distribution, structures, trends, productive and adaptive performances of populations of the existing breeds. As stated in preceding sections, although much information is lacking, conservation of farm AnGR in the Ethiopian perspective should be viewed from the rational utilization and protection of existing genotypes from genetic introgression. Unfortunately, no conservation activities of farm AnGR have so far been practised in the country, except for limited activities that are meant to maintain pure stocks of three cattle and one sheep breeds (Annex Table 5.1). Breeds of farm AnGR so maintained are Borena cattle breed at Did Tuyura Ranch, Horro cattle breed at Horro Ranch, Fogera cattle breed at Metekel Ranch and Adet Agricultural Research Centre, and Menz sheep breed at Amed Guya Research Centre.

1.7. The State of the Art in AnGR Management

In Ethiopia, information for sustainable utilization and conservation of the farm AnGR are very limited and, if available, are full of gaps.

1.7.1. Phenotypic characterization

In this document, phenotypic characterization refers to the morphological descriptions of farm AnGR only. Attempts by the MoA in 1975 were one of the few pioneering efforts made to classify the indigenous sheep, goat and cattle breeds into specific groups. Based on the tail form and quality of the hair, sheep were classified into four broad categories as Hairy Thin Tailed, Woolled Thin Tailed, Fat Tailed and Fat Rumped. Goats were classified into five major groups as Nubian, Highland, Afar, Somali and Long Tailed Gishe. Similarly, Alberro (1982a, 1982b) has classified cattle into four broad categories, namely: the Humpless Hametic Longhorn and Shorthorn, the Zebu, the Senga and the Intermediate Senga.

The other nationwide baseline survey was conducted on goats by FARM Africa in collaboration with the MoA and AUA (Alemayehu Reda, 1993; Workneh Ayalew, 1994; Nigatu Alemayehu, 1994).

A survey to initiate community-based conservation practices has also been conducted on the status of Borena cattle breed by BLPD/GTZ in collaboration with Oromia Regional Bureau of Agricultural Development (Alemayehu Reda, 2001). Oromia Agricultural Research Institute (Bako and Adami Tulu) is undertaking phenotypic characterization on sheep and goats. Amhara Region Agricultural Research Institute (ARARI) has undertaken phenotypic characterization on sheep breeds of the region in 2001. Similarly, Tigray Regional Agricultural and Natural Resources Development Bureau conducted a survey on livestock breeds of the region in 1997. More recently, the Oromia Agricultural Development Bureau in collaboration with ILRI (ILRI, 2004) conducted a comprehensive survey on livestock breeds of the region.

1.7.2. Evaluation of breeds and crosses

A number of organizations attempted to evaluate performances of some farm AnGR at different times.

- The MoA and the IAR/EARO have been intensively involved in performance evaluation activities of farm AnGR. These are cattle breeds such as the Borena, Fogera, Arsi and their crosses with different exotic breeds within its ranches.

- IAR/EARO and AUA have also been evaluating performances of such cattle breeds as Horro, Begayit, Fogera and Borena and their crosses with different exotic breeds. IAR/EARO (Debre Zeit and Holetta) has evaluated performances of different chicken breeds, Horro, Borena and Begayit cattle breeds and their crosses.
- Oromia Agricultural Research Institute (Bako and Adami Tulu) has conducted performance evaluation on Borena and Horro cattle breeds and their crosses as well as Horro sheep breeds and Rift Valley Family goats.
- AUA has also evaluated performances of Somali and Afar camel breeds, Somali goat breeds, and Borena and Begayit cattle breeds and their crosses.
- Debub University has undertaken performance studies on Somali goat breeds and their crosses.
- Debre Zeit, Holetta and Adami Tulu research centres have evaluated performances of donkeys found in the central highlands. Major emphasis was placed on assessing productive and reproductive performances of the breeds and their crosses under improved feeding and management conditions.

Mekonnen Haile-Mariam and Hailu Kassa (1995) have estimated genetic parameters for growth traits of Borena cattle breed. Similarly, Million Tadesse and Alemayehu Reda (2000) evaluated reproductive and productive performances of Borena cattle and their crosses with HF. Million Tadesse (2001) estimated crossbreeding parameters for milk production traits of Borena, Begayit and their crosses with HF. Million Tadesse and Tadelle Dessie (2003) have also evaluated performances of local, HF, and their crosses.

The Debre Zeit Agricultural Research Centre (Alemu Yami et al., 1998) has evaluated egg production and growth performances of indigenous chickens. Workneh Ayalew (2000) has evaluated performances of some well-known sheep and goat breeds.

1.7.3. Genetic characterization

In Ethiopia, only limited activities on genetic characterization have been conducted on some breeds of cattle, goats and chicken.

In the study conducted by Hanotte et al. (2000) on indigenous African ruminant livestock; specifically on cattle, sheep and goats, molecular characterization meant to quantify between-and within-population genetic diversity at microsatellite INRA124 was conducted on 10 cattle breeds namely: Bale, Borena, Ogaden, Sheko, Nuer, Raya-Azebo, Danakil, Horro, Arado and Fogera cattle breeds. Characterization of nine goat breeds, namely: Afar, Western Highland, Hararghe Highland, Western Lowland, Anglo-Nubian, Toggenberg breeds, crosses of Anglo-Nubian with Somali, Hararghe Highland and Toggenberg with Hararghe Highland have been conducted by Addis Ababa University (Addisu Simachew, 2002) using blood protein polymorphism. Genetic characterization of indigenous Tililli, Jarso, Cheffe, Tepi and Horro chicken breeds using Fayoumi (exotic) as reference has been conducted by Tadelle et al. (2003). Oromia Agricultural Research Institute has estimated genetic parameters of Horro sheep. Similarly, ARARI has estimated genetic parameters of Menz sheep breed and their crosses with Awassi.

PART TWO

2. CHANGING DEMANDS ON LIVESTOCK PRODUCTION AND THEIR IMPLICATIONS FOR FUTURE POLICIES, STRATEGIES AND PROGRAMMES

2.1. Impacts of Past and Present Policies, Strategies and Programs on AnGR Management

2.1.1. Government policies

Planned national economic development programs have been undertaken in Ethiopia since the early 1960's. Over the past five decades, the country has undergone through three distinct political changes along with corresponding food and agricultural policies that have influenced the development of the livestock sub-sector.

During the imperial regime up to 1974, livestock development interventions focused on creation of public institutions that support and promote improved livestock production, marketing, processing and distribution. Favourable government policies such as subsidized milk collection, secured land tenure, and free markets for feed and other inputs as well as economics of scale in production and marketing had encouraged large private commercial farming. The emergence of modern dairy and poultry operations using exotic and high-grade genotypes was evident in peri-urban areas. In addition, provision of government services was launched to support smallholders who had little land, capital and poor access to improved breeding animals.

During the socialist-oriented regime between 1974 and 1991, the economic policy was characterized by government intervention and monopoly in all economic sectors. Activities traditionally belonging to the private sector such as meat and by-products processing plants, state dairy and poultry farms, cattle and sheep breeding and multiplication ranches, and livestock export were run by the government. Government support for smallholders was largely nominal and, with the intention of promoting

medium- and large-scale operations, support was available only for peasant members of producer or service cooperatives that were based exclusively in rural areas. Actual government support was for unprofitable state farms, the production that had little impact on the development of the livestock sub-sector.

From 1991 to the present, Ethiopia has been undergoing through structural adjustment programs and market liberalization. Major policy reforms are being taken in the macro-economic framework and in the sectoral and cross-sectoral policies and strategies. The food and agricultural policy is elaborated in a comprehensive document: "Rural Development Policies, Strategies and Instruments (RDPSI)" issued by the Government in November 2001. According to RDPSI, agriculture is given the priority to lead the overall economic development and measures are being taken to create a supportive and favourable policy environment to promote the private sector and to transform agriculture from subsistence to small-scale farming. Several Government strategies and programs, poverty reduction and ensuring rapid economic development highlighted the importance of livestock in the reduction of rural poverty, attaining food security and sustainable economic development. Like for other economic sectors, decision-making power on animal resources planning and management, and livestock support services are decentralized to regional states and local level stakeholders. Beside favourable Government policy support, the prospective market opportunity for the smallholder livestock producers looks more promising than ever before.

2.1.2. Interventions in the livestock sector

A series of programs and projects aimed at increasing food and animal production which may affect management of farm animal genetic resources have been undertaken both at institutional and producers level. The overall objective was achieving increased livestock production and productivity, and thereby enhancing its contribution to the national economy. Major efforts have been directed toward removing the constraints facing livestock development and transferring new technologies to transform the

traditional sector into a modern livestock production. Genetic improvement of the indigenous animal resources is an integral component of most of the interventions.

Efforts to improve genetic potentials of local animals have resorted to crossbreeding with the "improved" exotic breeds. The first introduction of exotic cattle breeds took place in 1947 when the United Nations Relief and Rehabilitation Administration (UNRRA) donated 300 Friesian and Brown Swiss dairy cattle under the post Second World War Relief Program, which were used as nucleus herd at state, owned dairy farms. From mid-1970s onwards when involvement of the public sector dominated, state owned parastatal dairy and poultry farms, and cattle and sheep improvement and multiplication ranches served as central sources of breeding stocks for distribution to smallholders. Several government organizations were responsible in the implementation of these interventions with some financial and technical assistance from international, bi- or multi-lateral agencies. The MoA and the Ministry of State Farms Development (MSFD) have been the major government institutions engaged in importing, multiplying and distributing exotic, high grade and/or F₁ crossbred animals to smallholders. To a limited extent, the former Relief and Rehabilitation Commission (RRC), now Disaster Prevention and Preparedness Commission (DPPC) and NGOs have also been involved in the distribution of crossbred or local cattle, sheep, goats and chickens for restocking in areas affected by drought and displacement. Artificial insemination of cattle, introduced in late 1930's and further expanded with the establishment of the National Artificial Insemination Centre (NAIC) in 1981, has facilitated widespread crossbreeding of the local cattle with European breeds.

2.1.3. Impact on animal production and AnGR diversity

Despite substantial public investment for development of the livestock sub-sector, the resulting impact on the contribution of the sub-sector to the overall economic growth was unsatisfactory. The significant share of Ethiopia in East Africa's total output is attributable mainly to the size of the livestock population rather than increased productivity.

However, results of farm-level studies have shown that adopting crossbred cows and the associated package of improved feeding and management strategies could increase

milk production and household income of resource-poor smallholder mixed crop-livestock farmers (ILRI, 1997). Furthermore, adoption of crossbred cows brought about improved human health and nutritional status of all members of the household (Jemal Haider et. al., 2000; Shapiro et. al., 2000).

Given suitable government policy support, access to market and services, there is great potential for development of smallholder dairy scheme in peri-urban and urban areas (Staal and Shapiro, 1996). However, it could still be argued whether the gains attained were commendable when compared to the substantial investment involved in the genotype improvement undertakings, and recent hypothesis suggested that the economic benefits of crossbreeding may have been overestimated as non-market effects and environmental values have not been included in breed comparison studies (Workneh Ayalew, 2002).

There is little quantified information as to the impact of livestock interventions on the diversity of indigenous farm AnGR. The extent to which the exotic genotypes have diffused into the indigenous populations and the level of dilution is not objectively assessed. But, available estimates indicated that crossbred cattle make only 1% of the total cattle population of Ethiopia. It is a general understanding among professionals that indiscriminate distribution of exotic chicken might have caused rapid genetic dilution and erosion of local chicken types.

At farm level, the genotype of crossbred animals kept by producers is likely to include any percentage of foreign genetic material due to unsystematic breeding practices, unplanned mating schemes uncontrolled AI services and bull distribution. There is no controlled breeding in communal grazing areas where exotic bulls are introduced and allowed to indiscriminately mate with local cows. Such schemes resulted in a complex mixture of genotypes and threats to indigenous AnGR. In the absence of clearly defined breeding strategies, crossbreeding is bound to run the risk of diluting, erosion or loss of adaptation and loss of probably unique genetic identity of indigenous genetic resource.

Past research and development efforts ignored the importance of adapted indigenous farm AnGR due to a general belief that they are not productive and incapable of contributing to increased agricultural production.

2.1.4. Shortcomings of previous efforts

Political instability and/or inconsistency of development policies: weakness has always been in institutionalizing and setting commitments for the continuation of programmed undertakings. Experiences reflect that lasting acceptances, stability and further enriching and carrying on of an on-going programs are missing. Moreover, various livestock improvement programs and projects have not often taken the specific conditions of different production environments, market and agro-ecological conditions of the country into account. The need for favourably modified production environment such as improvements in the physical, management and institutional environments have been neglected and as a result these schemes have not been sustained after completion of the projects.

Lack of stakeholder participation: participation of stakeholders in the identification of problems, formulation of solutions, implementation and monitoring of interventions is very low.

Limitation of interventions in scope and scale: past activities often focused only on a couple of species (usually cattle and poultry), while other species of major significance to the poor were totally ignored.

2.1.5. Lessons learnt

1. F₁ crossbred cattle are adaptable and produce more than locals under modified management environments in the highlands
2. the importance of indigenous resources and knowledge as a basis for development
3. the need for modified production environment complementary to genetic improvement

4. the need for comprehensive and integrated national livestock development plan and breeding strategies based on agro-ecological zones, production systems and breed types
5. the need for demand-driven research

2.2. Future Demands and Trends

2.2.1. Demand for animal products

Demand for animal products in developing countries is projected to rise significantly and faster than in developed countries as a result of population growth, urbanisation and rising incomes. Greater consumption diversification or changes in food patterns are likely to occur in the developed countries, as a larger proportion of consumers require a more varied menu of a higher quality (Delgado et. al., 1999).

Output of meat and milk in Ethiopia is low and growth in productivity has been lagging behind population growth rates. As a consequence, the trend in per caput output of livestock products has been negative. Between 1995 and 2000, total milk and meat production increased by 2.6% and 1.4% per year, respectively. According to Genet Mengistu (2004) with an estimated annual growth rate of 2.7%, the population of Ethiopia is projected to increase from 63.5 million in 2000 to 84.0 million in 2010 and 106.0 million in 2020. Figure 2.1 illustrates the wide gap in per capita consumption of animal products between Ethiopia, Sub-Saharan Africa (SSA) and developed countries and signifies the link between population growth and declining per capita animal food availability. The picture for milk appears much worse than for meat. Proportion of per caput intake of calories (5.8%) and proteins (13.7%) as compared to 6.3% and 19.5% in SSA and 26.5% and 56.1% in developed countries is also relatively smaller (FAO, 2001).

Additionally, demand for livestock products is expected to increase as a result of urbanization. The proportion of urban population in Ethiopia increased from about 13.8 % in 1994 to 15.0% in 2004 and is expected to reach 19.9% in 2020. Accordingly, the rural population is expected to feed 21.2 million in 2020 compared to

about 9.0 million in 2000. While the overall population increase suggests that overall food supply has to increase greatly, higher urbanisation would mean a shift in dietary preferences towards higher quality food items such as meat, milk and eggs.

Figure 2.1. Average per capita milk and meat consumption in Ethiopia, Sub-Saharan Africa and developed countries, 1997-1999 (Source: FAO, 2001).

Prospects for economic growth may not appear as encouraging as expected in the near future to influence demand for animal products. However, market-oriented, demand driven opportunities for growth exist where population concentration is increasing. A study on urban demand for live sheep showed that sheep prices and household income, as well as socio-demographic factors, including household size and composition, significantly affect the likelihood of buying and expenditures on live sheep. Projections of live sheep demand and supply in Addis Ababa in 2010 and 2020 showed that sheep producers in Addis Ababa alone will be able to meet up to only 27% of the demand (Ehui et. al., 2000). Thus, the impact of population growth plus urbanisation will provide impetus for livestock development

2.2.2. Change in livestock production systems

A low growth rate of livestock output vis-à-vis high human population growth rate becomes one of the major causes of concern in Ethiopia. Comparison of livestock population with the small share of total livestock output and very low output per

animal shows that the nature of the livestock production systems is the least developed and indicates that an all out efforts have to be made to increase production and productivity.

Increased demand for livestock products may provide an engine for sustainable intensification of smallholder food and feed production systems. Changing factor combination of land, labour and capital and rising prices for livestock products are the starting point for livestock producers to invest in technologies that help achieve higher yields of livestock products and generate higher incomes. Crossbred animals can be seen as such a technology, provided that adequate infrastructure, extension and technical support services are established. Availability of sources of dairy genotypes was cited by Walshe et al. (1991), as an important factor in establishing market oriented dairying.

Livestock traditionally have been an important source of income for the rural poor. It is anticipated that expansion of livestock and livestock products offers one of the best opportunities for increased trade and economic integration between pastoralists and farmers. Several African countries can also be considered as potential export markets given the present food deficit and declining per capita meat consumption.

Improved livestock marketing is viewed as an important national development strategy to increase both rural incomes and foreign exchange. The share of foreign trade in livestock and livestock products to the overall export value of Ethiopia between 1999 and 2002 is shown in Table 2.1. Skins and hides in the form of pickled, wet-blue, and finished leather and leather products are the major exports. Ethiopian hides and skins have relatively high demands in Western industrialized nations and South East Asian countries. Next to skins and hides, goat meat and chilled mutton as well as canned meat products are the second important export commodities. Live sheep, goats and camels and to a limited extent beef cattle are exported mainly to Middle East and African countries.

The Government of Ethiopia has decided to integrate its economy into the Multilateral Trading System. Recently, Ethiopia has applied for World Trade Organization membership. The agreement on agriculture was designed to bring the agricultural sector under new set of multilateral trade rules and disciplines, which cover the three components of agricultural liberalization: market access, export subsidies and domestic support.

Table 2.1. Export values of livestock and livestock products (millions US\$), 1999-2002

Item	USD (000) in year			
	1999	2000	2001	2002
Live cattle	130	73	503	336
Live sheep and goat	866	1,256	NA	35
Other animals	111	3	4	182
Hides	1,271	3,953	16,795	3,948
Skins	25,430	38,200	56,895	53,956
Leather products	3,080	5,310	164	23
Bees wax	895	893	555	554
Dairy products	46	4	10	20
Meat and meat products	3,799	2,360	1,515	1,710
Value of total products exported	431,659	481,040	427,451	413,226
Share of livestock to total export value (%)	8.33	10.85	17.94	14.75

Source: National Bank of Ethiopia and Ethiopian Customs Authority

NA = Not Available

Sanitary problems and products quality have always constrained the export trade of Ethiopian products. With increased liberalisation of both foreign and domestic markets of most developing countries, import and export of meat and milk may be expected to increase at a rate faster than is currently observed. Product properties such as quality and safety, and increasingly the production methods, will influence the choice of products. Sanitary requirements of the international market and increased competition are expected to encourage quality products and diversification of products.

The presence of problems such as unavailability of inputs, inadequate research application and extension support, poor infrastructure, and inadequate technologies, leading to insufficient animal productivity will limit the rate of adoption of technology toward intensification and commercialisation of specialised livestock farming systems.

Urban and peri-urban areas offer most favourable opportunity for the adoption of technology for the development of intensified and specialized dairy and poultry

production systems. Smallholder farmers in the highlands who carry out dairy activities within a traditional mixed farming system prefer to maintain crossbred cattle and some cattle of local breeds and, provided that they have better access to inputs, market outlet and research and extension support, they are likely to adopt medium level input dairy production. Utilization of crop residues, increased production of manure ensuring soil fertility and diminishing the need to purchase feed will remain important in ensuring sustainable integration of crop and livestock production.

The majority of rural farmers and pastoralists cannot follow this path of intensification because of increasing transaction costs and poor marketing infrastructures, although adoption of crossbred cattle appears attractive. The different opportunities in low-input systems include diversification of livestock products through promotion of post-harvest technologies and small-scale processing units, providing incentives for organic products, and improving quality aspects of traditional products.

For pastoralists and agro-pastoralists systems, the main thrust will be to devise and implement disease monitoring and approved certification schemes for export of live animals and animal products. Animal identification associated with sanitary control will be very important.

National efforts to provide smallholders and pastoralists with breeding services to improve genetic potentials of animals could be regarded as a public good and as such warrants government support. Animal performance recordings will have best chances to be adopted in medium to large private livestock farms that have considerable scope for increasing productivity, but will require more inputs supply and advisory services. Research and development institutions mandated for the different aspects of breed improvement need to coordinate their tasks, establish common livestock recording systems and ensure the common use of information.

2.3. Alternative Strategies in the Utilization and Conservation of AnGR

Widespread poverty, hunger, malnutrition and deteriorating environmental conditions are the problems the facing. Therefore, the major concern for the agriculture sector is to respond to the demand-driven production, while ensuring sustainable management of the natural resources. Local genetic resources are crucial for sustainable production in such environmental and economic circumstances. In this regard, valuation of local breeds should consider their major contribution to risk management under conditions of environmental fluctuation. The future emphasis will have to concentrate on increased livestock productivity per animal and per unit of input, mainly land. These call for the improvement of livestock management techniques and exploitation of new possibilities to increase efficiency.

Breeding strategies for sustainable use of farm animal diversity are not currently available. On the other hand, there is a strong tradition of livestock keeping in pastoral and mixed farming systems. Traditional selection methods improve productivity only very slowly. In the past, the approach was able to keep pace with increasing food demands. In the future this will not be the case. Biotechnology offers tools that achieve rapid increases in productivity. Breeding strategies to improve efficiency of livestock production comprise selection schemes amongst populations, crossbreeding and biotechnological techniques to enhance the productive and reproductive performance. Therefore, improved AnGR management envisages sustained and balanced resource utilization and conservation by developing work linkages among national, regional and global initiatives. Thus, the overall objective of strategies on AnGR management should ensure better use, development and conservation of farm AnGR diversity.

2.4. Future National Policy for Utilization and Conservation of AnGR

The future breeding policy on use, development and conservation of AnGR should develop strategy, guidelines for farmers, researchers, extension workers, and on suitable breeds for the various agro-ecological zones. It should also include alternative breeding programs, regulations of import and export of genetic materials; characterization,

conservation and sustainable use of indigenous genetic resources and use of modern breeding technologies. Priority areas for research, development and the legal and institutional framework required for implementation should be identified. There is also urgent need to build more constructive approaches to breeding, ensuring that valuable local types are conserved. At the same time, a guided approach is required for specialised breed substitution and crossbreeding programs. Addressing the above issues will assist in basing the groundwork required for a genetic improvement work.

PART THREE

3. THE STATE OF NATIONAL CAPACITIES AND FUTURE CAPACITY BUILDING REQUIREMENTS

3.1. State of Capacity on Farm AnGRs

3.1.1. Current institutional capacity

The limited knowledge on Ethiopia's animal diversity indicates that no substantive measures have been taken to prevent the growing genetic erosion. The scope and depth of animal genetic resources utilization and conservation are still underdeveloped. Inadequate organizational structure, coordination facilities, scientific capacity, decision-making mechanisms are further compounded by the lack of a comprehensive policy and legislation. Thus, there is an urgent need for the formulation of a coordinated institutional capacity for animal genetic resources use, development and conservation programs.

The Federal Government of Ethiopia is delineating greater responsibility of conservation, development and sustainable utilization of natural resources to the regional governments limiting itself to public functions such as issuing policy, legal directives and playing regulatory role. It is, therefore necessary to create community awareness and build up their capacity for sustainable use of farm AnGR. Building capacity at the federal and regional levels through bilateral agreements, exchanging experience through workshops, training, and creating other linkages would be beneficial. At the moment, attempt is being undertaken to synchronize efforts between the various institutions of higher learning, research, development and conservation institutions, on the one hand and policy makers of the country at all levels on the other hand.

Table 3.1. Institutions involved in the livestock sector

	Extension	Research	Breeding	Education	Domestic animal diversity
International		ILRI	ILRI		ILRI, FAO GTZ, IDAD
Regional		ASARECA			
National	MoA	EARO	NAIC	Universities	EPA, IBC, EARO
Regional	Regional agricultural bureaus	Regional agricultural research institutes	Breeding ranches	agric. college & training centres	regional agricultural research institutes
Local	Wereda agricultural development offices	Farmers	Peasant Associations		Farmers

3.1.2. Policies and strategies on AnGR

Ethiopia has made several global and regional agreements regarding the conservation and utilization of biological diversity. By ratifying the UN Convention on Biological Diversity (CBD, 1992), specifically Ethiopia has committed itself to develop sustainable management of genetic resources. The convention necessitates initiation of measures at a national level designed for identifying and monitoring biodiversity, maintaining and sharing the resulting benefit arising out of the utilization of genetic resources, and integrating the conservation and sustainable use of biological resources into national decision making (FAO, 1998). It is, thus, a general acknowledgement that development of animal agriculture in Ethiopia must be based on sustainable utilization and conservation of farm animal genetic resources.

In line with this perspective, the Government of Ethiopia recognized the need to promote environmentally sustainable social and economic development of the country through protecting and preserving environmental values including biodiversity. Accordingly, Ethiopia has adopted policies and strategies that directly address biodiversity conservation. These are the Federal environmental policy (1997), the National biodiversity conservation and research policy (1998), and the Conservation strategy of Ethiopia (1997) and Regional conservation strategies specific to each regional state.

The draft policy and strategies for livestock utilization and conservation prepared by the MoA (1992) have not yet been approved. However, the EARO, the MoA, Alemaya University of Agriculture, the IBC and other relevant institutions and organizations are making efforts to formulate the national livestock breeding policy.

Currently, there is an understanding of the vital contribution of AnGR to the national economy. At present, the Federal and Regional States finance most of the livestock research activities. There are also some limited financial and technical supports from international and non-governmental organizations.

3.1.3. Policies and strategies favoring AnGR

- Constitution of FDRE
- Rural development policy and strategy
- Industrial development strategy
- Education policy
- Federal environment policy
- Biodiversity conservation and research policy, strategies and programs
- Conservation strategy of Ethiopia
- Conservation strategies of regional states
- Inclusion of AnGR issues as an integral part of several governmental organizations (MoA, EARO, HLI)

However, a breeding policy has not been adopted nor institutionalized.

3.2. Current Research and Development Status and Facilities

In spite of the vast potential resources in farm animal diversity, the current capacity of the country in animal genetic resources research is very limited both institutionally and in terms of qualified manpower. Therefore, building up a research capability is essential and a prerequisite to the utilization and conservation of genetic resources.

Limited capacities of basic research facilities are available at both Federal and Regional research Institutions. Molecular biology laboratory at IBC, Animal health

laboratory at NVI and Faculty of Veterinary Science at Addis Ababa University, Bedelle and Sebeta Animal Health Laboratories are some of research facilities available in the country. In area of Biotechnology there is MOET laboratory at the ILRI Debre Zeit Research Station. However, these facilities are not sufficient to undertake experiments needed in animal genetic resources.

Recently a national biotechnology strategy and action plan has been developed. Moreover, there is a plan to establish an Institute of Biotechnology at the Holetta Agricultural Research Centre.

Government organizations, primarily the MoA, the former MSFD and to a limited extent the then RRC, now Disaster Prevention and Preparedness Commission (DPPC), have all been involved in importing, multiplying and distributing exotic, high grade or crossbred animals to smallholder farmers.

The NAIC is responsible for production and distribution of semen as well as for importing bulls and semen that are used mainly for the dairy development. It also recruits bulls for semen production both from the exotics and the local cattle.

International or bilateral agencies (UNICEF, WFP, SIDA, FINIDA, FAO, ADB, GTZ, World Bank...) provide technical advice and some financial input to the development of livestock production. ILRI is also involved in some areas of livestock research activities.

3.3. Training Institutions

Institutions of higher learning, notably Alemaya University and Jimma Agricultural College are pioneers in starting research on farm animals. Currently, almost all these institutions and ATVET centres are involved in undertaking training on farm animals. The ATVET program, which provides all-round training to extension agents in almost all the regions of the country, is vital in transferring knowledge from research to classroom and the farming community.

3.4. Natural Resources Conservation and Management

The Federal Government of Ethiopia has taken an action with regard to the conservation of the country's biological resources by establishing institutions like the IBC and the Environmental Protection Authority to enforce policies on biodiversity conservation and protection of the environments.

In June 1998, the then IBCR, was established by decree and published the National Policy on Biodiversity Conservation and Research in the same year. Besides, Ethiopia has not only participated in the United Nation's Conference on Environment and Development (UNCED) held in Rio de Janeiro in 1992, but also endorsed and signed the Convention on Biological Diversity (CBD).

The Animal Genetic Resources Program of the IBC will undertake studies on livestock breed identification, characterization, conservation, breed evaluation and studies on the diversity and distribution of breeds in the country.

3.5. Future Capacity Requirements

Performance recording schemes are not in place, but are planned. Private large-commercial, smallholders and rural farmers shall be responsible for recording. Utilization of improved technologies and methodologies in breed evaluation will have a great impact on the use and development of AnGR. As the country is yet to establish genetic improvement programs, the opportunity exists to start at the state of the art level of technology. Development of reproductive biotechnological tools and genomic have been effectively developed over the last twenty years and is now routinely used on farm animals to quickly multiply desirable animals and to enhance genetic selection. There is a need to build capacity in this area.

PART FOUR

4. STRATEGIC PRIORITIES FOR ACTION

4.1. State of Development of Farm AnGR

In the low input system cattle, small ruminants, equines, camels and chicken are the most important farm AnGR. The entire equine populations as well as the majority of other farm AnGR that are maintained within the system are composed of indigenous breeds. On the other hand, in the medium input production system are the indigenous breeds of cattle, poultry, crossbreeds and pure exotic breeds of both species.

The majority of the livestock production of the country is traditionally run and natural resources based low input production system. Despite this, four species of the farm AnGR of the country, namely: cattle, sheep, chicken and goats are addressed in different development programs. Four of the seven widely used cattle breeds (Borena, Horro, Arsi, and Fogera) are used for pure and cross breeding programs. Similarly, such programs address the use of pure and crossbred chickens with the exotic broilers and layers. Two sheep breeds (Menz and Horro) and two goat breeds (Short and Long-eared Somalis) are addressed in the program. However, there exist no programs for the improvement of equines and camels (Annex Table 4.4).

4.2. Opportunities, Challenges and Priorities for Utilization and Conservation of Farm AnGR

4.2.1. Opportunities

Opportunities for better use and development of AnGR exist in mixed crop-livestock production system. The broad genetic variability, the ability of the indigenous breeds of cattle, sheep, goats, chicken and honeybees to survive under stressful environmental conditions including high disease incidence and poor nutrition are some of the opportunities for use and development of farm AnGR.

Studies undertaken so far show that indigenous sheep and goat breeds have good growth and reproduction performances. The large framed Fogera and Horro cattle breeds have reasonably satisfactory levels of milk yield. Exotic cattle for dairy production, small ruminants for meat and milk (goat) and chickens for egg and meat production can be used in low input production mixed crop-livestock system. A recent study on scavenging chicken (Fayoumi) indicated the existence of great potential for rural poverty reduction and protein supply.

Opportunity also exists in beekeeping. Ethiopia is endowed with immense potential of diversified millineries of plants. There is over 7,000 species of flowering plants, most of which are utilized by the honeybees. Besides, the country's variable climate, ample water resources and other conducive ecological factors have enabled the country to sustain a large number of honeybee colonies.

In the pastoral and agro-pastoral areas, there are also opportunities for better use and development of AnGR. Based on limited findings at research stations and government breeding ranches, the Borena cattle are typified as potential beef and dairy breed. Ogaden cattle breed has good potential for beef production. Black Head Somali sheep, Horro sheep and Somali goat breeds have good potential for mutton production. Sheko and Nuer breeds are considered to be trypanotolerant. The Afar/Danakil cattle are very resistant and well adapted to extremely harsh environmental conditions. Beef ranching will create good opportunity for supply of domestic and international markets. Ethiopia has a comparative advantage in producing beef for Middle East and some other countries.

Opportunity for better use and development of AnGR exists in the urban and peri-urban areas also. Dairy cattle, chicken and small ruminants are the most important species in the system. Commercial farms use pure exotic and crossbred dairy cattle and chicken. The system is characterized by the presence of agro-industrial by-products. It also has good market opportunities for livestock and livestock products than the other production systems. The Existence of various research, teaching and development

institutions as well as other facilities in and around these areas makes the system more favourable for livestock production.

The indigenous knowledge involved in the traditional livestock production could be considered as an opportunity for the use and development of AnGR. Women and children play important roles in management and utilization of AnGR, particularly small ruminant and chickens. Women are also involved in feeding, watering and health management of large ruminants. Moreover, they play important roles in processing and marketing of livestock products. Children participate actively in milking and herding activities.

4.2.2. Challenges

Use, development and conservation of AnGR have been constrained by different challenges. One of the major challenges is weak institutional set up & capacity which resulted in lack of coordination between stakeholders, absence of responsible body at national level, inadequate institutions engaged in AnGR; low involvement of professional associations, civil societies, private sector and absence of breeders associations.

Inadequate human resources capacity, limitations in know-how and expertise, is the other major challenge for better use of AnGR. There are very few animal breeders, experts in modern reproductive technology and molecular genetics. The Ethiopian government is sensitised and committed to provide necessary support. However, the country is faced with the scarcity of financial resources. Therefore, additional financial support is required from national, regional and global initiatives, and the international community.

Unimproved management of natural and human resources have also contributed a lot to the under-development of AnGR.

Poor inputs and services delivery systems are the other challenges for sustainable use, development and conservation of AnGR. Inputs and services delivery systems are often fragmented, piece meal approach with no inter-links among development projects. Dairy development projects, for example, often focussed on high inputs delivery approach, such as distribution of crossbred in-calf heifers, which requires intensive delivery of AI and animal health services.

The performance of breed improvement and multiplication centres in supplying improved stock to the farmers has been low. The current dissemination scheme of crossbred, high grade and exotic animals, assisted by subsidized AI service, is unable to satisfy the ever-increasing demand by smallholders. In addition, there are a number of other technical, technological and socio-economic constraints that hinder improvements in productivity and efficiency of smallholders.

The other challenge in the use and development of AnGR is lack of coordination in development, research, training, extension and delivery of inputs. Efforts have not been institutionalised and tended to be short-term oriented.

Some of the major challenges of beef production are lack of proper understanding of the production systems, improper use natural resources, inadequate nutrition, fragile environment, prevalence of diseases, poor processing and marketing infrastructures.

Free and unregulated movement of livestock in the country has also contributed to uncontrolled breeding and dissemination of diseases.

Other challenges for better utilization and conservation of farm AnGR are poverty, recurrent drought and financial constraints.

Information and database management systems necessary for AnGR use, development and conservation are not available. There exist no comprehensive data on farm AnGR of the country. The limited available information on some cattle breeds is documented in the FAO (DAD-IS) and ILRI (DAGR-IS) databases.

The absence of livestock breeding policy, strategies and action plans has led to the uncontrolled interbreeding and crossbreeding of a number of species of animals, thus affecting the use, development and conservation of AnGR. This has also an impact on potential cooperation with other countries and international organizations.

4.2.3. Priorities

Developing institutional set up such as strengthening of coordination between stakeholders, capacitating of institutions engaged in AnGR; promoting of the roles of professional associations, civil societies and private sector, and formation of breeders associations is one of the major priorities.

To implement AnGR utilization and conservation programs, there is a need for capacity building in animal breeding, reproduction, nutrition, health and other related disciplines. Therefore, development of human resource in areas of characterization, utilization and conservation are the other priority areas.

Effective research and development programs geared towards better utilization and conservation of AnGR require strong collaboration and networking with various national, regional and international institutions. Similarly, formation of breed societies and breeders associations will have an important role in AnGR use, development and conservation.

There is a need to develop database management system, information network and early warning system at a national level to capture national, regional and global information on use, development and conservation of farm AnGR. In addition, establishment of a national recording system is one of the priorities. There is a need to establish systems to monitor the status and trends of breeds.

Classification, description, and identification of breeds of farm animals, including genetic and molecular level characterization breed level census and breeds bio-geographic distribution, is one of the top priority areas for better utilization and

conservation of AnGR. As a matter of urgency, there is a need to conserve the already endangered breeds such as the Sheko cattle breed.

PARTS FIVE AND SIX

5. THE WAY AHEAD: MAINSTREAMING ANIMAL GENETIC RESOURCES

5.1. Capacity Building

5.1.1. Communication between regions and countries

Establishment of a strong communication system and linkages between regions, regional institutions and countries that are engaged in utilization and conservation of AnGR is one of the options to build national capacity in AnGR utilization, development and conservation. For example, networking with the Biotechnology Trust for Africa (BTA), the Kenyan Agricultural Research Institute (KARI) and ASARECA is essential.

5.1.2. Capacity building

Capacity building requirements are:

- Establishment of a critical mass of a core team of national experts to form the basis for better use, development and conservation of AnGR, especially in the areas of genetics, breeding, genomics, biotechnology, research and development
- Provision of various levels of training to all stakeholders that are directly or indirectly involved in the use, development and conservation of AnGR
- Arranging short-term consultants to assist in the use, development and conservation of AnGR
- Setting up a well-equipped modern laboratory use, development and conservation of AnGR
- Formation of breed societies and breeder associations. They will have an important role to play in the use, development and conservation of AnGR
- Creation of institutional arrangements, which could effectively mobilise all stakeholders involved in AnGR activities.

5.1.3. Support to the continuity of regional workshops on AnGRs

Regional workshops are important vehicles to share experiences and identify gaps and challenges among countries in the region. Therefore, strong interaction between the National Focal Points (NFP) and support from FAO and other relevant institutions is required to implement such activities.

5.1.4. Develop regional markets for products from AnGR

Strong and sustainable national, regional and international markets should be developed to ensure sustainable use, development and conservation of AnGR. To this end, due attention should be given to the development of necessary infrastructure and institutions. In order to be competent in marketing of animals and animal products, establishment of disease free zones, standards and quality control are required.

5.1.5. Raising awareness

Awareness of farmers, general public and decision-makers on the strategic relevance of AnGR should be raised. To do so, establishment of a strong national database on AnGR and use of mass media are required. To increase awareness on the importance of AnGR, FAO and professional associations should lobby governments and policy makers.

Dissemination of indigenous knowledge in animal production and management of AnGR is highly important. To appreciate and develop the knowledge, strong extension service, commitment of stakeholders and finance are required.

5.1.6. Set up of gene banks

Strengthening the capacity of IBC and other relevant institutions that are dealing with use, development and conservation of AnGR need be supported by the government, FAO and other stakeholders.

5.2. Institutional Arrangements

For better utilization and conservation of AnGR, a well-coordinated institutional arrangement is a key issue. The arrangement should include dissemination of information and networking at regional and international levels. To implement this, concerned national, regional and international institutions and NFP should take up the lead responsibility. FAO is expected to play a catalytic role in this regard.

5.3. Possible Sources of Finance

The government, the private sector, communities, NGOs, FAO, bi-lateral and multi-lateral donors, World Association of Animal Production and other donors are potential sources of finance to activities geared towards better utilization and conservation of AnGR.

5.4. Policy Development

Policies favouring the promotion of use, development and conservation of farm AnGR should be developed in collaboration with the stakeholders.

5.5. Partnership

It is important to create and develop partnerships with national, regional and international institutions that are directly or indirectly involved in the utilization and conservation of AnGR.

5.6. Potential Modalities for Facilitating Interaction Among Donors and Recipients

Projects/proposals should be developed in the area of utilization and conservation of AnGR by stakeholders for funding. NCC should be a responsible body to coordinate these activities.

5.7. Process Forward: Post-State of the World Process

The NCC should further be strengthened and work closely with the concerned national, regional and international institutions. The NCC should play active role in lobbying government bodies to raise awareness on the importance AnGR. FAO should participate in influencing governments.

REFERENCES

- Addisu Simachew (2002). Characterization of some goat populations in Ethiopia by means of blood protein polymorphism. MSc Thesis, Addis Abeba University, Ethiopia.
- Alberro, M. and Solomon Haile-Mariam (1982a). The indigenous cattle of Ethiopia, Part I. *World Anim.Rev.*41: 2-10.
- Alberro, M. and Solomon Haile-Mariam (1982b). The indigenous cattle of Ethiopia, Part II. *World Anim. Rev.*41: 27-42.
- Alemayehu Reda (1993). Characterization (phenotypic) of indigenous goats and goat husbandry practices in East and South-eastern Ethiopia. MSc. Thesis, Alemaya University of Agriculture, Ethiopia, 135 pp.
- Alemayehu Reda (2001). Borena rangeland pastoral development program/GTZ (BLPDP/GTZ), Australian Agricultural Consulting and Management Company (1984). *Livestock Sub-sector Review, Volume I, Annex 3.*
- Alemu Yami and Tadelle Dessie (1998). The status of poultry research and development in Ethiopia. *Proceedings of the 5th Annual Conference of the Ethiopian Society of Animal Production (ESAP), Addis Ababa, Ethiopia, pp.40-60.*
- Amhara Regional Agricultural Development Bureau (2001). *Annual report on the status of livestock development of Amhara region.*
- Ayalew, W., King, J.M., Bruns, E. & Rischkowsky, B. (2003). Economic evaluation of smallholder subsistence livestock production: lessons from an Ethiopian goat development program. *Ecological Economics* 45 (3): 331-339.
- Azage Tegegne, Million Tadesse, Alemu Yami and Yoseph Mekasha (2000). Characterisation of urban and peri-urban dairy production in Ethiopia. *Urban Agriculture Magazine*, 1(2): 23-24.
- Bekele Tafesse and Tezera Getahun (1998). Preliminary observation on camel types and major parasitic diseases of camel in the Eastern lowlands of Ethiopia. *Proceeding of the 6th Annual Conference of the Ethiopian Society of Animal Production (ESAP)(, Addis Ababa, Ethiopia, pp. 201-207.*
- Bench Maji Agricultural Development Office (2001). *Annual report on livestock population of the Bench Maji Zone.*

- Beyene Kebede and Beruk Yemane (1992). Animal genetic resources and breed characterization work in Ethiopia. In: African Animal Genetic Resources: their characterization, conservation and utilization, Proceedings of the Research Planning Workshop held at ILCA, Addis Ababa, Ethiopia, 19-21 Feb., 1992.
- Bureau of Agriculture and Natural Resources Development of Tigray Regional State (1997). Tigray Regional State Livestock Development Action Program 1997. 110pp.
- Central Statistics Authority (2003). Ethiopian Agricultural Sample Enumeration for the year 2001/2002, Statistical report on farm management practices, livestock and farm implements, Part II, Addis Ababa, Ethiopia.
- Coppock D. L (1994). The Borena Plateau of Southern Ethiopia. Synthesis of pastoral research, development and changes 1980-1991, International Livestock Center for Africa (ILCA), Addis Ababa, Ethiopia.
- Delgado, C., M. Rosegrant, H. Steinfeld, S. Ehui, and C. Courbois (1999). Livestock to 2020 The Next Food Revolution. Food, Agriculture and the Environment Discussion Paper 28.
- Ethiopian Agricultural Research Organization (2002). Livestock research methodology: options for economic growth and to enhance the livelihoods smallholder farmers. A Paper presented to workshop on Poverty Reduction through Transforming Smallholder System from Subsistence to Market Orientation, June 3-7, 2003, Addis Ababa, Ethiopia.
- Ehui S.K., Benin S. and Nega Gebreselassie (2000). Factors affecting urban demand for live sheep: The case of Addis Ababa, Ethiopia. Socio-economics and Policy Research Working Paper 31. ILRI (International Livestock Research Institute), Nairobi, Kenya. 32 pp.
- Food and Agriculture Organization (1999). Animal genetic resources information No. 25. Rome, Italy.
- Food and Agriculture Organization (200). Preparation of the First Report on the State of the World's Animal Genetic Resources. 3. Guidelines for the development of country reports (unpublished), Rome, Italy.
- Food and Agriculture Organization (2001). FAOSTAT. FAO statistical databases on agriculture, fisheries, forestry and nutrition, Rome, Italy.

- FAO (2001). FAOSTAT. The State of the World's Plant Genetic Resources for Food and Agriculture, Rome, Italy.
- FARM Africa (1996). Goat types of Ethiopia and Eritrea: physical description and management systems. International Livestock Research Institute, Ethiopia, 55pp.
- Genet Mengistu (2004). Population and development in Ethiopia. A Paper presented at Women's Forum ACP-EU Joint Parliamentary Assembly, Feb. 14, 2004, UN CON. Center, Addis Abeba, Ethiopia.
- Getachew Asmenew, Ercole Zerbini and Abate Tedla (1993). Crop-livestock interactions and implications for animal traction research in the Ethiopian highlands. In: Proceedings of Fourth National Livestock Improvement Conference, 13-15 Nov., 1991, Institute of Agricultural Research, Addis Abeba, Ethiopia, pp. 29-36.
- Getachew Feleke (2003). Assessments of the Ethiopian Dairy sub-sector, Addis Abeba, Ethiopia.
- Hanotte O., Tawah C.L, Bradley D.G., Okomo M., Verjee Y., Ochieng J. and Rege J.E. O. (2000). Geographic distribution and frequency of a Taurine *Bos taurus* and an Indicine *indicus* Y specific allele amongst sub-Saharan Africa cattle breeds. *Molecular Ecology*, vol. 9; pp 387-396.
- International Livestock Research Institute (1997). Livestock, people and the environment International Livestock Research Institute, Nairobi, Kenya.
- International Livestock Research Institute (2004). Description, execution and analysis of livestock breed survey in Oromia Regional State, ILRI, Addis Abeba, Ethiopia.
- Jemal Haider, Shapiro B.I., Tsegaye Demisse, Alemu G/Wold (2000). The nutritional and health status of women and children in households with and without crossbred cows in Holetta Wereda, Ethiopia. In: Jabbar M.A., Peden D.G., Mohamed Saleem M.A. and Li Pun H. (eds), *Agro-ecosystem, natural resources management and human health related research in East Africa. Proceedings of an IDRC–ILRI international workshop held at ILRI, Addis*

- Ababa, Ethiopia, 11–15 May 1998. ILRI (International Livestock Research Institute), Nairobi, Kenya. pp. 124–136.
- Mekonnen Haile-Mariam and Hailu Kassa Mersha (1995). Estimates of direct and maternal (co) variance components of growth traits in Boran cattle. *J. Anim. Breed. Genet.* 112: pp 43-52.
- Million Tadesse (2001). Estimation of crossbreeding parameters for milk production traits of dairy cows at Debre Zeit Agricultural Research Centre, Ethiopia. *Ethiopian Journal of Animal Production* 1(1): 45-54.
- Million Tadesse and Alemayehu Reda (2000). Evaluation of reproduction and productivity of local Boran cows and Boran x Friesian heifers at Abernosa Cattle Breeding Ranch. Proceedings of the 8th Annual Conference of the Ethiopian Society of Animal Production (ESAP), Addis Ababa, Ethiopia, Pp. 206-210.
- Million Tadesse and Tadelles Dessie (2002). Evaluation of crossbreeding parameters of local Arsi and Holstein Friesian crosses at Assela cattle breeding centre. *Ethiopian Journal of Animal Production (EJAP)* 3: 25-35.
- Million Tadesse, Amha Kassahun and Alemu Regassa (2002). Domestic animal biodiversity strategy and action plan, draft document. In: Biodiversity Strategy Action Plan, IBCR, Addis Ababa, Ethiopia.
- Million Tadesse and Tadelles Dessie (2003). Evaluation of performance of local, Holstein Friesian and their crosses at Debre Zeit Research Centre, Ethiopia. *Livestock Research for Rural Development* 15:1-9.
- Ministry of Agriculture (2003). Beef Production Pilot Project, draft document, Addis Ababa, Ethiopia.
- Ministry of Agriculture (2004). Memorandum on the foreign trade regime: Ethiopia, main draft document (text), Addis Abeba, Ethiopia.
- National Conservation Secretariat, Ministry of Natural Resources Development and Environmental Protection (1994). Policy on the resource base, its utilization and planning for sustainability, Transitional Government of Ethiopia, vol.1, Addis Ababa, Ethiopia.

- Nigatu Alemayehu (1994). Characterization of indigenous goat types of Eritrea. Northern and Western Ethiopia. MSc. Thesis, Alemaya University of Agriculture, Alemaya, Ethiopia. 136pp.
- Regional Government of Tigray, Bureau of Agriculture and Natural Resources Development (1997). Tigray Livestock Development Action Program, pp 110.
- Shapiro, B.I., Jemal Haider, Alemu G/Wold and Abebe Misgina (2000). The intra-household economic and nutritional impacts of market-oriented dairy production: Evidence from the Ethiopian highlands: In: Agro-ecosystems, natural resources management and human health related research in East Africa, Jabbar, M..A. D.G. Peden, M.A. Mohamed Saleem and H. Li Pun (eds), proceedings of an IDRC–ILRI international workshop held at ILRI, Addis Ababa, Ethiopia, 11–15 May 1998, ILRI (International Livestock Research Institute).
- Sileshi Ashinie (2003). Managing water for livestock and fisheries development. In: Integrated water and land management research and capacity building priorities for Ethiopia. Proceedings of a MoWR/EARO/IWMI/ILRI international workshop, ILRI, Addis Abeba, Ethiopia, 2-4 Dec. 2002.
- Solomon Haile-Mariam (1981). Animal health review in Ethiopia (1972-1979), MoA, Livestock Resources Development Dept., Vet. Service Division, Addis Ababa, Ethiopia.
- Staal S., Delgado C. and Nicholson C. (1997). Smallholder dairying and transactions costs in East Africa. *World Development* 25:779-794.
- Tadelle Dessie (2003). Phenotypic and genetic characterization of local chicken ecotypes in Ethiopia, Dissertation, Landwirtschaftlich-Gaertnerischen Fakultät der Humboldt-Universität zu Berlin, Germany, ISBN 3-89574-497-2, 208pp.
- Teketel Forssido (1986). Studies on the meat production potential of some local strains of chickens in Ethiopia, PhD Thesis, J.L University of Giessen, 210pp.
- Walshe, M.J., Grindle, J., Nell, A., and Bachmann, M. (1991). Dairy development in sub-Saharan Africa: a study of issues and options. World Bank Technical Paper No 135. Africa Technical Department Series, pp.97.
- Wilson R.T. (1984). The camel in Ethiopia and Eritrea, Longman, London.

Workneh Ayalew (2000). Do smallholder farmers benefit more from crossbred (Somali x Anglo-Nubian) than from indigenous goats? Dissertation, University of Goettingen, Cuvillier Verlag, Goettingen, Germany, ISBN 3-89873-007-7.

Workneh Ayalew, Rey B., Peacock C.P. and Kano Banjaw (1994). Identification of goat breed types in Southern Ethiopia, Small Ruminant Research. In: Goat Types of Ethiopia and Eritrea. Physical description and management systems. 55pp.

Yishak Mengesha, S. Ehui, M. Jabbar and B. Shapiro (1998). Livestock production, consumption and trade: Key indicators. Livestock Policy Analysis Brief No. 11. International Livestock Research Institute (ILRI), Addis Ababa, Ethiopia.